



Spatial Distribution and Abundance of Culicodes Imicola and Obsoletus Group in Europe

Cuellar, Ana Carolina; Skovgård, Henrik; Nielsen, Søren Achim; Stockmarr, Anders; Anderson, G.; Lindström, Anders; Chicrico, J ; Lilja, T.; Lühken, R.; Steinke, S.

Total number of authors:
35

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Cuellar, A. C., Skovgård, H., Nielsen, S. A., Stockmarr, A., Anderson, G., Lindström, A., Chicrico, J., Lilja, T., Lühken, R., Steinke, S., Kiel, E., Larska, M., Hamnes, S. I., Sviland, S., Hopp, P., Brugger, K., Rubel, F., Balenghien, T., Garros, C., ... Bødker, R. (2016). *Spatial Distribution and Abundance of Culicodes Imicola and Obsoletus Group in Europe*. Abstract from 3rd Conference on Neglected Vectors and Vector-Borne Diseases , Zaragoza, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

SPATIAL DISTRIBUTION AND ABUNDANCE OF *CULICOIDES IMICOLA* AND *OBSOLETUS* GROUP IN EUROPE

Cuellar A.C.¹, Skovgaard H.², Nielsen S.A.³, Stockmarr A.⁴, Anderson G.⁵, Lindstrom A.⁵, Chirico J.⁵, Lilja T.⁵, Lühken R.⁶, Steinke S.⁶, Kiel E.⁶, Larska M.⁷, Hamnes S.I.⁸, Sviland S.⁸, Hopp P.⁸, Brugger K.⁹, Rubel F.⁹, Balenghien T.¹⁰, Garros C.¹⁰, Rakotoarivony I.¹⁰, Allène X.¹⁰, Lhoir J.¹⁰, Delécolle J.C.¹¹, Mathieu B.¹¹, Delécolle D.¹¹, Setier-Rio M.L.¹², Venail R.¹², Scheid B.¹², Miranda Chueca M.A.¹³, Barcelo Segui C.¹³, Lucientes J.¹⁴, Estrada R.¹⁴, Wesley Tack¹⁵, Mathis A.¹⁶ and Bødker R.¹

¹Section of Epidemiology, Technical University of Denmark (DTU)-National Veterinary Institute, Copenhagen, Denmark

²Department of Agroecology - Entomology and Plant Pathology, Aarhus University, Aarhus, Denmark

³Institute for Science and Environment. Roskilde University, Roskilde, Denmark

⁴Department of Applied Mathematics and Computer Science, Technical University of Denmark (DTU), Copenhagen, Denmark

⁵National Veterinary Institute (SVA), Uppsala, Sweden

⁶Bernhard Nocht Institute for Tropical Medicine, WHO Collaborating Centre for Arbovirus and Haemorrhagic Fever Reference and Research National Reference Centre for Tropical Infectious Diseases, Hamburg, Germany

⁷Department of Virology, National Veterinary Research Institute, Pulawy, Poland

⁸Norwegian Veterinary Institute, Oslo, Norway

⁹Institute for Veterinary Public Health, Vienna, Austria

¹⁰Agricultural research for development (CIRAD), Montpellier, France

¹¹Institute of parasitology and tropical diseases of Strasbourg, Université de Strasbourg, Strasbourg, France

¹²EID Méditerranée, Montpellier, France

¹³Laboratory of Zoology. University of the Balearic Islands. Spain

¹⁴Department of Animal Pathology. University of Zaragoza. Spain

¹⁵Avia- GIS. Zoersel, Belgium

¹⁶Institute of Parasitology, University of Zürich, Zürich, Switzerland

Corresponding author: anacu@vet.dtu.dk

Introduction

Biting midges (Diptera, Ceratopogonidae) of the genus *Culicoides* are important vectors of veterinary important pathogens causing large economic losses in many European countries. *Culicoides imicola* and the *Obsoletus* group are considered to be the main vectors of bluetongue virus in Europe that affects cattle and sheep. Spatial-temporal modelling of the vector distribution

and abundance allows to identify areas under risk of virus transmission and can help to apply effective surveillance and control measurements.

Materials and Methods:

For this purpose, presence-absence and abundance data of *Culicoides* were collated from 1005 sites across 9 countries (Spain, France, Denmark, Poland, Switzerland, Austria, Poland, Sweden, Norway) between the years 2007 and 2013. The dataset included information on the vector species abundance (number of specimens caught per night), GPS coordinates of each trap, start and end dates of trapping. A dataset of 120 environmental variables obtained from a Fourier transformed MODIS temporal series were used with the Random Forest machine learning algorithm in VecMap developed by AviaGIS to predict the overall species distribution and monthly abundance in Europe.

Results:

For every month of the year, maps were generated to visualize the abundance of *C. imicola* and *Obsoletus* group in Europe and a distribution map for each of both showing the probability of occurrence. Although the variance was large, the predicted abundance values for each site had a positive correlation with the observed abundance.

Spatial variation was found in abundance for *C. imicola* and the *Obsoletus* group. For *C. imicola* the highest abundance was found in Spain as for the *Obsoletus* group the highest abundance was found in eastern France. Temporal variation was also observed with the highest abundance occurring during summer months and the lowest during winter months

Conclusion

We were able to model the spatial distribution in Europe for *C. imicola* and *Obsoletus* group in terms of abundance and suitability. The maps obtained showed a consistent pattern with the reported distribution for *C. imicola* and the *Obsoletus* group. The observed seasonal variation was also consistent with the reported by literature as *Culicoides* population dynamic depends on environmental factors such as temperature and rainfall. The outputs obtained here can be used as inputs for epidemiological models and can be helpful for determining areas under risk of transmission.